

GRAZERS

**Grazers contains calcium and other ingredients:
calcium chloride solution -
10% calcium oxide, 7.2% w/w calcium**
Should not be mixed with any other product

Application rate

Basic rate: 250ml in 100-250 litres of water per ha. **Knapsack rate:** 20ml in 20 litres water.

A double rate can be safely applied if damage persists.

Application timing

Apply to *dry foliage* (avoid frost and strong sunlight) at any stage of growth during the growing season (very effective on first true leaf), as soon as the damage has occurred, or to areas potentially at risk.

GRAZERS is rainfast in one hour and can remain effective for up to 6 weeks, but in certain situations, where crops are grazed, mowed, irrigated or exposed to prolonged heavy rainfall after spraying, repeat applications may be required sooner.

Grazers is very safe to use and beneficial to the plant.

Arable (Cereals, Oil Seed Rape, Root Crops, Game Cover)

Grazers can be applied safely and effectively to any crop at risk from damage.

Cost effective treatments

Rabbits: a headland spray (min. 24m boom) is generally sufficient unless severe pest numbers.

Pigeons: treat areas at risk (eg. under powerlines, below roosts) at early leaf emergence, and weaker areas during late winter to stimulate re growth and allow crop to recover.

Note: with waxy leaved crops eg. brassicae it can assist if an inert wetter UN65 is added or just a small amount of household detergent.

Horticulture (Salad/Veg, Ornamental Crops, Grass)

Grazers can be applied safely and effectively to any crop at risk from damage.

N.B. Modules, liners & transplants are extremely vulnerable.

Grazers is specifically designed to stimulate growth of the crop beyond the susceptible stages for grazing damage.

Spray at least twice at double strength before planting/putting out, allow the leaf to dry between applications. Spray again 2-3 days later in the field.

Note: with waxy leaved crops eg. brassicae it can assist if an inert wetter UN65 is added or just a small amount of household detergent.

Precautions

Avoid contact with skin and eyes. Ingestion may cause vomiting. If irritation or symptoms persist seek medical advice.

Grazers **must** be sprayed on its own a minimum of 3 days before or after any other spray application.